

How to use multiply marked *multicent* strains for mapping genes and translocation breakpoints.

David D. Perkins

Background

Normal-sequence linkage testers have been constructed that contain readily scored markers near the centromeres of all seven linkage groups (Table 1; Perkins 1972, 1990). While these *multicent* testers can be used for assigning unmapped genes to linkage group, their main value in our hands has been to identify the chromosomes involved in new centromere-linked translocations, which are recognized because they result in linkage between markers that are normally independent. The first choice for mapping new genes is usually *alcoy*; *csp-2* (Perkins and Björkman 1979), which has fewer and more easily scored markers than those in the *multicent* testers. However, because three unlinked translocations are already present in *alcoy*; *csp-2*, crosses may fail to reveal the linkages of a new, unmapped translocation that involves one of the same chromosome pairs that is already translocated in the tester.

One disadvantage with the original *multicent-1* tester was presence of a colonial marker *bal* (*balloon*) which interfered with scoring other markers except in *bal*⁺ progeny. Modified testers without balloon were therefore constructed and designated *multicent-3* to *-5* (Table 1). (Metzenberg *et al.*, 1984, had independently created *multicent-2* for use in RFLP mapping.)

Procedure

Heterokaryons of *multicent* strains with the inactive-mating-type *helper-1* strain *a^{m1} ad-3B cyh-1* (FGSC 4564; Perkins 1984) are phenotypically wild type and are fertile as female parents on unsupplemented crossing medium. (The *helper-1* component does not participate in the cross. See *How to use alcoy for linkage group assignment*.)

The *multicent-3* tester incorporates two changes: *arg-5* is substituted for *bal* in linkage group II, and a long inversion, *OY323*, is inserted as a crossover-suppressor in linkage group I. Progeny are isolated either to complete medium or to minimal supplemented with arginine and pyridoxine. As a result of the heterozygous inversion, *mating type* shows little or no recombination with markers throughout two-thirds of LG I, the longest linkage group.

The *multicent-4* tester also has *arg-5* substituted for *bal*. In addition, *psi* replaces *pdx-1* as a marker for IV. *psi* (*protein-synthesis-inhibited*) is a readily-scored temperature-sensitive conditional mutant that does not grow or survive at 34°C although it is normal on minimal medium at 25°. *arg-5* is scored by transferring progeny to minimal medium at 25°C, *psi* by transfer to arginine-supplemented minimal at 34°. Ascospores must be germinated at 25°C.

multicent-5 differs from *multicent-4* only in having the *OY323* inversion present in linkage group I.

Scoring for the other markers is as follows: Tests for mating type are most readily accomplished on *fluffy* lawns in petri dishes (Perkins *et al.* 1989. See *How to determine mating type*). The *at* mutant (*attenuated* morphology) is readily scorable on minimal (with or without supplements) at 2 or 3 days (34°C). Growth is flat on the surface, with scattered specks of conidiation. In *wc-1* (*white collar-1*), carotenoids are absent in mycelia but not in conidia. Scoring *wc-1* is better at 34°C than at 25°.

Germinants are incubated until conidia become orange, preferably under illumination. Scoring of *ylo-1* (*yellow-1*) is unreliable in young (3- or 4-day) cultures, especially in combination with *wc-1*, but it becomes increasingly clear with age. The color difference is more apparent under some light sources than

others, so if difficulty is experienced distinguishing *ylo* from the orange wild type, different fluorescent or other sources of illumination should be tried. *acr-2* is clearly resistant to 50 µg/ml acriflavine on solid medium. Small inocula should be used when scoring *pdx-1*.

With any of the testers, progeny are scored for markers sequentially, beginning with the visible markers *at*, *wc-1*, and *ylo-1*. If linkage is apparent, the remaining markers need not be tested. If the unmapped mutant is unlinked to a visible marker, the markers that require transfer are then tested serially until linkage is seen. With translocations, the normally independent *multicent* markers are examined for linkages to one another. Linkage, of course, is indicated if parental combinations are significantly more frequent than recombinants among the progeny. For ratios deviating significantly from 1:1, see table in Perkins (1944) or in *How to use genetic methods for detecting linkage*.

The new *multicent* testers are heterokaryon-compatible with strains of OR background (*het-C het-d; het-e; het-x^{OR}*). They are available as highly fertile phenotypically wild-type heterokaryons in combination with the inactive-mating-type *helper-1* strain (Table 1). Although the homokaryotic testers can be used for crossing, labor can be minimized by using a helper-assisted heterokaryon as female parent, improving fertility and making it unnecessary to supplement the crossing medium.

Figure 1, which is a tally sheet for scoring markers in crosses by *multicent-4*, can be adapted for use with other *multicent* testers.

References

Metzenberg, R. L., J. N. Stevens, E. U. Selker, and E. Morzycka-Wroblewska. 1984. A method for finding the genetic map position of cloned DNA fragments. *Neurospora Newslett.* 31: 35-39.

Perkins, D. D. 1972. Linkage testers having markers near the centromere. *Neurospora Newslett.* 19: 33.

Perkins, D. D. 1984. Advantages of using the inactive-mating-type *a^{ml}* strain as a helper component in heterokaryons. *Neurospora Newslett.* 31: 41-42.

Perkins, D. D. 1990. New *multicent* linkage testers for centromere-linked genes and rearrangements in *Neurospora*. *Fungal Genet. Newslett.* 37: 31-32.

Perkins, D. D. 1991. *Neurospora alcoy* linkage tester stocks with group VII marked, and their use for mapping translocations. *Fungal Genet. Newslett.* 38: 83.

Perkins, D. D. 1994. Deviations from 1:1 and numbers of progeny necessary for establishing linkage. *Fungal Genet. Newslett.* 41: 69-70.

Perkins, D. D., B. C. Turner, V. C. Pollard, and A. Fairfield. 1989. *Neurospora* strains incorporating *fluffy*, and their use as testers. *Fungal Genet. Newslett.* 36: 64-67.

DDP

Table 1. Constitution of multivalent testers 1 through 5

Strain designation	I	II	III	IV	V	VI	VII
<i>multivalent-1</i>	<i>mat A</i> or <i>a</i>	<i>bal</i>	<i>acr-2</i>	<i>pdx-1</i>	<i>at</i>	<i>ylo-1</i>	<i>wc-1</i>
<i>multivalent-2</i>	<i>mat a un-2</i>	<i>arg-5</i>	<i>thi-4</i>	<i>pyr-1</i>	<i>lys-1 inl</i>	---	<i>nic-3 ars-1</i>
<i>multivalent-3</i>	<i>In(IL;IR)OY323</i> <i>mat A</i> or <i>a</i>	<i>arg-5</i>	<i>acr-2</i>	<i>pdx-1</i>	<i>at</i>	<i>ylo-1</i>	<i>wc-1</i>
<i>multivalent-4</i>	<i>mat A</i> or <i>a</i>	<i>arg-5</i>	<i>acr-2</i>	<i>psi</i>	<i>at</i>	<i>ylo-1</i>	<i>wc-1</i>
<i>multivalent-5</i>	<i>In(IL;IR)OY323</i> <i>mat A</i> or <i>a</i>	<i>arg-5</i>	<i>acr-2</i>	<i>psi</i>	<i>at</i>	<i>ylo-1</i>	<i>wc-1</i>

FGSC Numbers

Strain designation	<i>A</i>	<i>a</i>	<i>A</i> + <i>helper-1</i>	<i>a</i> + <i>helper-1</i>			
<i>multivalent-1</i>	2014	2015					
<i>multivalent-2</i>		4488					
<i>multivalent-3</i>	6824	6825	6826	6827			
<i>multivalent-4</i>	6828	6829	6830	6831			
<i>multivalent-5</i>	6832	6833	6834	6835			

Tally for Multicentromere Tester 4

Cross No. _____ Strain or locus tested: _____

Parents: arg-5; acr-2^R; psi; at; ylo-1; wc-1; _____ X _____

Linkage of unknown to markers (un = _____)

			at-un	wc-un	ylo-un	acr-un	arg-un	psi-un	mt-un
(P)	+	-							
(P)	-	+							
(R)	+	+				(P)			
(R)	-	-				(P)			

Figure 1.

How to use *multicent* strains for mapping genes and translocation breakpoints.