

Chromosomal mapping of genes encoding subunits of complex I from *Neurospora crassa*.

Fátima Ferreira, Teresa Almeida, Margarida Duarte and Arnaldo Videira - Instituto de Biologia Molecular e Celular, Unidade Multidisciplinar de Investigação Biomédica and Instituto de Ciências Biomédicas de Abel Salazar, University of Porto, Portugal.

The genes encoding the 21 kDa protein and the 19.3 kDa, 21.3 kDa and 24 kDa iron-sulfur subunits of complex I were located, by RFLP analysis, in Linkage Groups IV, VI, VI and V, respectively, of the *Neurospora crassa* genome.

Complex I or respiratory chain NADH dehydrogenase is located in the inner mitochondrial membrane, where it catalyzes electron transfer from NADH to ubiquinone through a series of protein-bound prosthetic groups, coupled to proton translocation across the membrane. In mammals and *N. crassa*, seven subunits of this mitochondrial enzyme are encoded and synthesized in the organelle, whereas about 30 others are imported from the cytoplasm. The nuclear-coded genes are usually called *nuo* (for NADH:ubiquinone oxidoreductase) followed by the molecular mass of the respective polypeptide. We have determined the chromosomal location of four new nuclear genes encoding subunits of complex I by the analysis of segregation of RFLPs among the strains of the multicent-2 cross kit (Metzenberg *et al.* 1984 Fungal Genet. Newsl. 31:35-39). Using cDNAs encoding proteins of 19.3 kDa, 21 kDa, 21.3 kDa and 24 kDa as probes (the homologues of bovine PSST, AQDQ, TYKY and 24 kDa proteins, respectively; Azevedo *et al.* 1994 Biochim. Biophys. Acta 1188:159-161; Duarte *et al.* 1996 Biochim. Biophys. Acta 1275:151-153; unpublished results), we have isolated the respective genomic clones from a λ J1 library obtained from the FGSC. Genomic DNA fragments containing the genes were then labeled with different non-radioactive techniques employing either PCR or random-primer labeling, and used in the RFLP analysis. The genes encoding the 21 kDa protein and the 19.3 kDa, 21.3 kDa and 24 kDa iron-sulfur proteins were located in Linkage Groups IV, VI, VI and V, respectively, of the *N. crassa* genome (Table 1). With these assignments, ten nuclear-coded single-copy genes encoding subunits of complex I have been located (reviewed in Videira 1998 Biochim. Biophys. Acta, in press).

Table 1. Segregation of RFLPs, related to complex I genes, among the strains of the multicent-2 cross kit

	AABBCCDDEEEFFGGHHIIJJKKLLMMNNOOPPQQRR
	14671457135713145768141414582324142414
<i>nuo19.3</i>	M-MOMMMMOOMMMMOOOOOO-OOOMMOOOOMMMMOOMM
<i>nuo21</i>	MOMMMOOMMMOOMOMOMOOOM-OOMOMOMMOM-OMMMMO
<i>nuo21.3c</i>	M-OOMOOMOMOMOMMOOOOMOOOOMOMOMOMMOMOMOOM
<i>nuo24</i>	M-O-MOOMOOMMMOOO-MOO-MMOOM-MOMOMMMOMM-